arXiv:0812.1984v1 [astro-ph] 10 Dec 2008

Potential-density pairs for a family of finite disks

Earl Schulz

60 Mountain Road, North Granby, CT 06060

`earlschulz@gmail.com`

ABSTRACT

Exact analytical solutions are given for the three finite disks with surface density $\Sigma_n = \sigma_0(1 - R^2/\alpha^2)^{n-1/2}$ with $n = 0, 1, 2$. Closed-form solutions in cylindrical coordinates are given using only elementary functions for the potential and for the gravitational field of each of the disks.

The n=0 disk is the flattened homeoid for which $\Sigma_{hom} = \sigma_0/\sqrt{1 - R^2/\alpha^2}$. Improved results are presented for this disk. The n=1 disk is the Maclaurin disk for which $\Sigma_{Mac} = \sigma_0\sqrt{1 - R^2/\alpha^2}$. The Maclaurin disk is a limiting case of the Maclaurin spheroid. The potential of the Maclaurin disk is found here by integrating the potential of the n=0 disk over α, exploiting the linearity of Poisson's equation. The n=2 disk has the surface density $\Sigma_{D2} = \sigma_0\left(1 - R^2/\alpha^2\right)^{3/2}$. The potential is found by integrating the potential of the n=1 disk.

Subject headings: gravitation- methods: analytical-celestial mechanics- galaxies: kinematics and dynamics - galaxies: individual(NGC 891)

1. Introduction

Observations of edge-on galaxies now provide detailed structural and kinematic 3-D information. However attempts to use this data to generate mass distribution have not been completely successful. One problem is that it is difficult to calculate the potential and force vectors for finite disks. There is a need for fully solved finite disks which can be used for theoretical studies, for benchmarking computer programs, and as basis functions to directly model 3-D observations.

Mass modeling commonly assumes that the disk mass in contained in an infinite disk. These include the exponential disk (Freeman 1970), the Mestel disk (Mestel 1963), the Kuzmin-Toomre disk (Toomre 1963; Binney & Tremaine 2008; Evans & de Zeeuw 1992; Conway 2000), and the Rybicki disk (Evans & Collett 1993).

Only a few finite disks have been solved analytically for all (R, z). Lass & Blitzer (1983) and Vokrouhlicky & Karas (1998) give a solution for a for all (R, z) for a finite thin disk with constant density. The gravitational attraction approaches infinity at the edge of this disk. The gravitational attraction of the finite Mestel disk (Mestel 1963; Lynden-Bell & Pineault 1978; Hunter et al. 1984) and the truncated exponential disk (Casertano 1983) are finite at the edge of the disk but these disks have not been solved in closed form for points off the disk. Huré et al. (2008); Huré (2005) describe a method to approximate the potential of a power law disk for points both on and off the disk.

The family of finite disks with surface density $\Sigma_n(R;\alpha) = \left(1 - R^2/\alpha^2\right)^{n-1/2}$ are related to the Maclaurin spheroid which has been studied since the time of Newton. This family has recently been studied by González & Reina (2006) and Pedraza et al. (2008). González & Reina (2006) use the method of Hunter (1963) to obtain the general solution as a sum of Legendre polynomials in elliptical coordinates and give evaluated expressions for the potentials for the disks 1, 2, and 3. Here we derive complete closed form solutions in cylindrical co-ordinates for the potential and the gravitational fields of the n=0, 1, and 2 disks. We simplify the integration by moving to the imaginary domain in way which is similar to the complex-shift method introduced by Appell. See Ciotti & Marinacci (2008); Ciotti & Giampieri (2007) and references therein

The gravitational potential follows Poisson's equation $\nabla^2\Phi = 4\pi \mathrm{G}\rho$. Poissons's equation is linear so that solutions may be added. That is, if $\nabla^2\Phi_1 = 4\pi \mathrm{G}\rho_1$ and $\nabla^2\Phi_2 = 4\pi \mathrm{G}\rho_2$ then $\nabla^2(\Phi_1+\Phi_2) = 4\pi \mathrm{G}(\rho_1+\rho_2)$. Similarly, solutions may be differentiated with respect to a parameter to obtain a new solution. Toomre (1963) and Lynden-Bell (1989) exploited this linearity to find a family of velocity-density pairs by differentiating the Kuzmin (1956) model . In the same way, Satoh (1980) derived new models from the Miyamoto & Nagai (1975) set of potential-density pairs by differentiating with respect to a parameter. Here we use a similar approach by integrating a known solution with respect to a parameter. If the integral is tractable the result is a new potential-density pair.

Maple 11 was used in this work. Maple was invaluable in simplifying the unwieldy intermediate results but needed a good deal of coaxing to handle the messier equations. The Maple results were cross checked in several ways.

A common notation is used throughout. Cylindrical co-ordinates (R, z) are used; Σ is the disk surface density; σ_0 is the surface density at $R = 0$; and α is the disk radius. Φ is the potential where Φ is always negative and $\Phi(\infty) = 0$; F_R and F_z are the gravitational field vectors with the standard sign convention, ie: $\boldsymbol{F} = -\boldsymbol{\nabla}\Phi$.

The principal values of elementary functions are used so that, for instance, for $z = x + \mathtt{I}\,y = r\cos(\theta) + \mathtt{I}\,r\sin(\theta)$; $\sqrt{z} = \sqrt{r}\cos(\theta/2) + \mathtt{I}\,\sqrt{r}\sin(\theta/2)$, valid for $-\pi < \theta < \pi$. In this way the function $\sqrt{z}$ is unambiguous with continuous derivatives except on the negative real axis where it is discontinuous.

2. The n=0 Disk: The Flattened Homeoidal Shell

A homeoid is a shell of uniform density which is bounded by similar spheroids. Newton was the first to prove that the net force is 0 (i.e., the potential is constant) within these shells. See Chandrasekhar (1987) for historical background. The infinitely thin homeoidal shell is a differential element of a spheroid. The homeoidal disk is the limiting case for which the minor axis approaches 0. The collapsed density of the thin homeoid is:

$$\Sigma_{hom}(R;\alpha) = \begin{cases} \sigma_0/\sqrt{1-R^2/\alpha^2} & \text{for } R<\alpha \\ 0 & \text{for } R>\alpha \end{cases} \tag{1}$$

Lynden-Bell (1989) gives a formula for the external potential of the thin homeoidal shell. Taking the limit $c = (1-e^2)^{1/2}\alpha \rightarrow 0$ gives a solution for the disk which is valid at all R and z.

Cuddeford (1993) gives an expression for the potential of this disk which is simpler than previous solutions:

$$\Phi_{hom}(R,z;\alpha) = -2\pi\alpha\sigma_0\mathrm{G}\sin^{-1}\left[\frac{2\alpha}{\sqrt{z^2+(R+\alpha)^2}+\sqrt{z^2+(R-\alpha)^2}}\right] \tag{2}$$

Equation 2 can be further simplified. First make the trivial transformation:

$$\Phi_{hom}(R,z;\alpha) = -2\pi\alpha\sigma_0\mathrm{G}\sin^{-1}\left[\frac{\sqrt{z^2+(R+\alpha)^2}-\sqrt{z^2+(R-\alpha)^2}}{2R}\right] \tag{3}$$

Now use the identity A1 to obtain

$$\Phi_{hom}(R,z;\alpha) = -\pi\alpha\sigma_0\mathrm{G}\ \left[\sin^{-1}\left(\frac{\alpha-\mathtt{I}\,z}{R}\right)+\sin^{-1}\left(\frac{\alpha+\mathtt{I}\,z}{R}\right)\right] \tag{4}$$

Equation 4 can be integrated over α whereas Equation 3 yields an impossible integral. Equation 4 must be real valued on physical grounds. This is easy to prove by noting that $\sin^{-1}(\mathrm{x})$ is an odd function of x and so the odd powers of $\mathtt{I}\,z$ in the series expansion of equation 4 cancel, leaving a real valued result.

The gravitational field for the collapsed homeoid is obtained from the potential. Equation 4 yields particularly simple expressions for the field vectors $F_{R,hom}$ and $F_{z,hom}$:

$$F_{R,hom}(R,z;\alpha) = -\frac{\pi\alpha\sigma_0\mathrm{G}}{R}\left[\frac{\alpha-\mathtt{I}\,z}{\sqrt{R^2-(\alpha-\mathtt{I}\,z)^2}}+\frac{\alpha+\mathtt{I}\,z}{\sqrt{R^2+(\alpha-\mathtt{I}\,z)^2}}\right] \tag{5}$$

$$F_{z,hom}(R,z;\alpha) = -\pi\alpha\sigma_0\mathrm{G}\left[\frac{\mathtt{I}}{\sqrt{R^2-(\alpha-\mathtt{I}\,z)^2}}-\frac{\mathtt{I}}{\sqrt{R^2-(\alpha+\mathtt{I}\,z)^2}}\right] \tag{6}$$

These force vectors can be expressed as entirely real functions using identities A8 and A9:

$$F_{R,hom}(R,z;\alpha) = -\sqrt{2}\pi\alpha\sigma_0 \mathrm{G}\, \frac{\alpha\sqrt{f_1 f_2 - f_3} - \mid z \mid \sqrt{f_1 f_2 + f_3}}{R f_1 f_2} \tag{7}$$

$$F_{z,hom}(R,z;\alpha) = -\sqrt{2}\pi\alpha\sigma_0 \mathrm{G}\, \frac{\mathrm{sgn}(z)\sqrt{f_1 f_2 + f_3}}{f_1 f_2} \tag{8}$$

Where

$$\begin{aligned} f_1 &= \sqrt{z^2 + (R+\alpha)^2} \\ f_2 &= \sqrt{z^2 + (R-\alpha)^2} \\ f_3 &= \alpha^2 - R^2 - z^2 \end{aligned} \tag{9}$$

3. The n=1 Disk: The Maclaurin disk

Beginning in the early 18'th century Colin Maclaurin, along with James Ivory and many others, studied the properties of elliptical bodies. Chandrasekhar (1987) includes a very good historical summary. See also Binney & Tremaine (2008); Bertin (2000); Schmidt (1956); Mihalas & Routly (1968); Kalnajs (1971, 1972).

The homogeneous oblate spheroid is the simplest case of a spinning body for which the gravitational attraction balances the centrifugal force. The Maclaurin disk, also known as the Kalnajs disk (Kalnajs 1972), is a limiting case for which minor axis is 0. The Maclaurin disk is defined by the surface density:

$$\Sigma_{Mac}(R;\alpha) = \begin{cases} \sigma_0\sqrt{1 - R^2/\alpha^2} & \text{for } R < \alpha \\ 0 & \text{for } R > \alpha \end{cases} \tag{10}$$

There are a few solutions for the potential of the Maclaurin disk in the literature. Mihalas & Routly (1968) gives an expression for the potential of an oblate homogeneous spheroid based on the derivation in Schmidt (1956). The potential of the Maclaurin disk can be found by letting the eccentricity $e \to 1$ while holding the mass constant. Hunter (1963) gives the solution for the Maclaurin disk as a series of Legendre polynomials in elliptical coordinates. Neugebauer & Meinel (1995); Meinel (2001); González & Reina (2006) give a closed form solution for the potential of the Maclaurin disk in elliptic coordinates.

The starting point here is the potential-density pair for the n=0 disk, the flattened homeoid for which $\Sigma_{hom}(R;\alpha) = \sigma_0/\sqrt{1 - R^2/\alpha^2}$. The surface mass density of the Maclaurin disk is found from the transformation:

$$\Sigma_{Mac}(R;\alpha) = \frac{1}{\alpha}\int_0^\alpha \Sigma_{hom}(R;\hat{\alpha})\, d\hat{\alpha} = \frac{1}{\alpha}\int_0^\alpha \frac{\sigma_0}{\sqrt{1 - R^2/\hat{\alpha}^2}}\, d\hat{\alpha} = \sigma_0\sqrt{1 - R^2/\alpha^2} \tag{11}$$

The corresponding potential is:

$$\Phi_{Mac}(R,z;\alpha) = \frac{1}{\alpha}\int_0^\alpha \Phi_{hom}(R;\hat\alpha)\; d\hat\alpha = \frac{-\pi\sigma_0 \mathrm{G}}{\alpha}\int_0^\alpha \hat\alpha\left[\sin^{-1}\left(\frac{\hat\alpha - \mathtt{I}\,z}{R}\right) + \sin^{-1}\left(\frac{\hat\alpha + \mathtt{I}\,z}{R}\right)\right] d\hat\alpha \tag{12}$$

Where the expression for Φ_{hom} is given by equation 4 above. Use integral 2.813 and 2.833 from Gradshteyn & Ryzhik (1994) to obtain:

$$\begin{aligned}\Phi_{Mac}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{4\alpha}&\Bigg[(2\alpha^2 - R^2 + 2z^2)\left(\sin^{-1}\left(\frac{\alpha + \mathtt{I}\,z}{R}\right) + \sin^{-1}\left(\frac{\alpha - \mathtt{I}\,z}{R}\right)\right)\\ &+ \alpha\left(\sqrt{R^2 - (\alpha - \mathtt{I}\,z)^2} + \sqrt{R^2 - (\alpha + \mathtt{I}\,z)^2}\right)\\ &- 3\;z\left(\mathtt{I}\sqrt{R^2 - (\alpha - \mathtt{I}\,z)^2} - \mathtt{I}\sqrt{R^2 - (\alpha + \mathtt{I}\,z)^2}\right)\Bigg]\end{aligned} \tag{13}$$

Equation 13 can be converted to an entirely real expression by using the identities A1, A6 and A7:

$$\begin{aligned}\Phi_{Mac}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{4\alpha}&\Bigg[2(2\alpha^2 - R^2 + 2z^2)\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ \sqrt{2}\alpha\sqrt{f_1 f_2 - f_3} - 3\sqrt{2}\mid z\mid\sqrt{f_1 f_2 + f_3}\Bigg]\end{aligned} \tag{14}$$

where f_1, f_2, f_3 are given by equation 9 above.

The gravitational field of the Maclaurin disk is found from the potential using Φ as given by equation 13. The resulting expressions were expressed as entirely real functions by using the identities A1, A8, and A9 :

$$\begin{aligned}F_{R,Mac}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{2R\alpha}&\Bigg[2R^2\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ \sqrt{2}\alpha(\alpha^2 - R^2 + z^2)\frac{\sqrt{f_1 f_2 - f_3}}{f_1 f_2}\\ &- \sqrt{2}\mid z\mid(\alpha^2 + R^2 + z^2)\frac{\sqrt{f_1 f_2 + f_3}}{f_1 f_2}\Bigg]\end{aligned} \tag{15}$$

$$\begin{aligned}F_{z,Mac}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{\alpha}&\Bigg[-2z\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ 2\sqrt{2}\alpha z\frac{\sqrt{f_1 f_2 - f_3}}{f_1 f_2}\\ &+ \sqrt{2}\;\mathrm{sgn}(z)\;(\alpha^2 - R^2 - z^2)\frac{\sqrt{f_1 f_2 + f_3}}{f_1 f_2}\Bigg]\end{aligned} \tag{16}$$

where f_1, f_2, f_3 are given by equation 9 above.

The potential on the z axis and on the $z = 0$ plane can be found by by taking limits of equation 14:

$$\Phi_{Mac}(0, z; \alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{\alpha}\left[\left(\alpha^2 + z^2\right)\sin^{-1}\left(\frac{\alpha}{\sqrt{\alpha^2 + z^2}}\right) - \alpha \mid z \mid\ \right] \tag{17}$$

$$\Phi_{Mac}(R, 0; \alpha) = \begin{cases} -\frac{\pi^2\sigma_0 \mathrm{G}}{4\alpha}\left(2\alpha^2 - R^2\right) & \text{for} R \leq \alpha \\ -\frac{\pi\sigma_0 \mathrm{G}}{2\alpha}\left[(2\alpha^2 - R^2)\sin^{-1}(\frac{\alpha}{R}) + \alpha\sqrt{R^2 - \alpha^2}\right] & \text{for} R \geq \alpha \end{cases} \tag{18}$$

The radial force vector in the $z = 0$ plane is found by taking the limit of equation 15 or by differentiating equation 18 with respect to R.

$$F_{R,Mac}(R, 0; \alpha) = \begin{cases} -\frac{\pi^2 R\sigma_0 \mathrm{G}}{2\alpha} & \text{for} R \leq \alpha \\ -\frac{\pi\sigma_0 \mathrm{G}}{\alpha}\left[R\sin^{-1}(\alpha/R) - \alpha\sqrt{1 - \alpha^2/R^2}\right] & \text{for} R \geq \alpha \end{cases} \tag{19}$$

The axial force vector on the z axis is found by taking the limit of equation 15 or differentiating equation 17 with respect to z.

$$F_{z,Mac}(0, z; \alpha) = -\frac{2\pi\sigma_0 \mathrm{G}}{\alpha}\left[z\sin^{-1}\left(\frac{\alpha}{\sqrt{\alpha^2 + z^2}}\right) - \alpha\ \mathrm{sgn}(z)\right] \tag{20}$$

4. The n=2 Disk

González & Reina (2006) give a closed form solution for the potential of the n=2 disk in elliptic coordinates.

The disk surface density of the n=2 disk is

$$\Sigma_{D2}(R; \alpha) = \begin{cases} \sigma_0(1 - R^2/\alpha^2)^{3/2} & \text{for } R < \alpha \\ 0 & \text{for } R > \alpha \end{cases} \tag{21}$$

This mass distribution can be obtained from equation 10, the disk surface density of the n=1 disk, with the transformation:

$$\Sigma_{D2}(R; \alpha) = \frac{3}{\alpha^3}\int_0^{\alpha} \hat{\alpha}^2 \Sigma_{Mac}(R; \hat{\alpha}) d\hat{\alpha} \tag{22}$$

The corresponding potential is:

$$\Phi_{D2}(R, z; \alpha) = \frac{3}{\alpha^3}\int_0^{\alpha} \hat{\alpha}^2 \Phi_{Mac}(R, z; \hat{\alpha})\ d\hat{\alpha} \tag{23}$$

Where the expression for Φ_{Mac} is given by equation 13 above. Equation 23 contains terms which after change of variable have with the form $\int x^n \sin^{-1}(x)dx$ and $\int x^n\sqrt{1 \pm x^2}dx$ and can be solved using Gradshteyn & Ryzhik (1994) 2.262, 2.813, and 2.833. After collecting terms, the result is reasonably compact:

$$\begin{aligned}\Phi_{D2}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{64\alpha^3}\Bigg[&3(8\alpha^4 - 8\alpha^2R^2 + 16\alpha^2z^2 + 3R^4 - 24R^2z^2 + 8z^4)\left(\sin^{-1}\left(\frac{\alpha + \mathtt{I}\,z}{R}\right) + \sin^{-1}\left(\frac{\alpha - \mathtt{I}\,z}{R}\right)\right)\\ &+ \alpha(18\alpha^2 - 9R^2 + 26z^2)\left(\sqrt{R^2 - (\alpha + \mathtt{I}\,z)^2} + \sqrt{R^2 - (\alpha - \mathtt{I}\,z)^2}\right)\\ &- z(58\alpha^2 - 55R^2 + 50z^2)\left(\mathtt{I}\,\sqrt{R^2 - (\alpha + \mathtt{I}\,z)^2} - \mathtt{I}\,\sqrt{R^2 - (\alpha - \mathtt{I}\,z)^2}\right)\Bigg]\end{aligned} \tag{24}$$

Equation 24 can be converted to an entirely real expression by using the identities A1, A6 and A7:

$$\begin{aligned}\Phi_{D2}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{64\alpha^3}\Bigg[&6(8\alpha^4 - 8\alpha^2R^2 + 16\alpha^2z^2 + 3R^4 - 24R^2z^2 + 8z^4)\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ \sqrt{2}\alpha(18\alpha^2 - 9R^2 + 26z^2)\sqrt{f_1f_2 - f_3}\\ &- \sqrt{2}\mid z\mid(58\alpha^2 - 55R^2 + 50z^2)\sqrt{f_1f_2 + f_3}\Bigg]\end{aligned} \tag{25}$$

where f_1, f_2, f_3 are given by equation 9 above.

The gravitational field of the n=2 disk is the gradient of the potential using Φ as given by equation 24. The resulting expressions were expressed as entirely real functions by using the identities by using identities A1, A8, and A9 :

$$\begin{aligned}F_{R,D2}(R,z;\alpha) = -\frac{3\pi\sigma_0 \mathrm{G}}{16R\alpha^3}\Bigg[&2R^2(4\alpha^2R^2 - 3R^4 + 12R^2z^2)\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ \sqrt{2}\alpha(2\alpha^4 - 5\alpha^2R^2 + 4\alpha^2z^2 + 3R^4 - 25R^2z^2 + 2z^4)\frac{\sqrt{f_1f_2 - f_3}}{f_1f_2}\\ &+ \sqrt{2}\mid z\mid(2\alpha^4 + 9\alpha^2R^2 + 4\alpha^2z^2 - 13R^4 - 11R^2z^2 + 2z^4)\frac{\sqrt{f_1f_2 + f_3}}{f_1f_2}\Bigg]\end{aligned} \tag{26}$$

$$\begin{aligned}F_{z,D2}(R,z;\alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{4\alpha^3}\Bigg[&- 6z(2\alpha^2 - 3R^2 + 2z^2)\sin^{-1}\left(\frac{f_1 - f_2}{2R}\right)\\ &+ \sqrt{2}\alpha z(13\alpha^2 - 13R^2 + 17z^2)\frac{\sqrt{f_1f_2 - f_3}}{f_1f_2}\\ &+ \sqrt{2}\ \mathrm{sgn}(z)((4\alpha^4 - 8\alpha^2R^2 - 3\alpha^2z^2 + 4R^4 - 7R^2z^2 - 11z^4)\frac{\sqrt{f_1f_2 + f_3}}{f_1f_2}\Bigg]\end{aligned} \tag{27}$$

where f_1, f_2, f_3 are given by equation 9 above.

The potential on the z axis and on the $z = 0$ plane can be found by by taking the limit of equation 25:

$$\Phi_{D2}(0, z; \alpha) = -\frac{\pi\sigma_0 \mathrm{G}}{4\alpha^3}\left[3\left(\alpha^4 + 2\alpha^2 z^2 + z^4\right)\sin^{-1}\left(\frac{\alpha}{\sqrt{\alpha^2 + z^2}}\right) - \alpha \mid z \mid (5\alpha^2 + 3z^2)\right] \tag{28}$$

$$\Phi_{D2}(R, 0; \alpha) = \begin{cases} -\frac{3\pi^2\sigma_0\mathrm{G}}{64\alpha^3}\left(8\alpha^4 - 8\alpha^2 R^2 + 3R^4\right) & \mathrm{for} R \leq \alpha \\ -\frac{3\pi\sigma_0\mathrm{G}}{32\alpha}\left[(8\alpha^4 - 8\alpha^2 R^2 + 3R^4)\mathrm{sin}^{-1}(\frac{\alpha}{R}) + 3\alpha(2\alpha^2 - R^2)\sqrt{R^2 - \alpha^2}\right] & \mathrm{for} R \geq \alpha \end{cases} \tag{29}$$

The radial force vector in the $z = 0$ plane is found by taking the limit of equation 26 or differentiating equation 29 with respect to R.

$$F_{R,D2}(R, 0; \alpha) = \begin{cases} -\frac{3\pi^2 R\sigma_0\mathrm{G}}{16\alpha^3}(4\alpha^2 - 3R^2) & \mathrm{for} R \leq \alpha \\ -\frac{3\pi\sigma_0\mathrm{G}}{8\alpha^3}\left[R(4\alpha^2 - 3R^2)\mathrm{sin}^{-1}(\alpha/R) - \alpha(2\alpha^2 - 3R^2)\sqrt{1 - \alpha^2/R^2}\right] & \mathrm{for} R \geq \alpha \end{cases} \tag{30}$$

The axial force vector on the z axis is found by taking the limit of equation 27 or differentiating equation 28 with respect to z.

$$\begin{aligned} F_{z,D2}(0, z; \alpha) = -\frac{\pi\sigma_0\mathrm{G}}{2\alpha^3(\alpha^2 + z^2)}\Bigg[&- 6(\alpha^2 + z^2)^2 z\mathrm{sin}^{-1}\left(\frac{\alpha}{\sqrt{\alpha^2 + z^2}}\right) \\ &+ \alpha z^2(13\alpha^2 + 17z^2) + \alpha\,\mathrm{sgn}(z)(4\alpha^4 - 3\alpha^2 z^2 - 11z^4)\Bigg] \end{aligned} \tag{31}$$

4.1. Comparison of the Maclaurin disk and the n=2 disk

Table 1 compares important properties of the two disks. Figure 1a compares the surface mass density and potential. Figure 1b compares rotational velocity in the disks. The n=2 disk is more centrally concentrated than the Maclaurin disk. The rotational velocity increases more quickly in the inner disk and begins to fall before reaching the edge of the disk. As is apparent from figure 1b, the derivative of the circular velocity of the Maclaurin disk is discontinuous at the edge of the disk whereas the n=2 disk is better behaved.



Fig. 1.— Compare the surface mass density and circular velocity, $V_c = \sqrt{(-VF_R(R,z)}$, for the Maclaurin disk and the n=2 disk. Reduced units are used so that the masses of both disks and the point mass are 1.0

Table 1 comparison of the Maclaurin and the n=2 disks

Property		**Maclaurin disk**	**n=2 Disk**
Surface density	$\Sigma(R) =$	$\sigma_0\sqrt{1 - R^2/\alpha^2}$	$= \sigma_0(1 - R^2/\alpha^2)^{3/2}$
Total mass	$M =$	$\frac{2}{3}\pi\alpha^2\sigma_0$	$= \frac{2}{5}\pi\alpha^2\sigma_0$
Circular velocity	$V_c^2(R,0) =$	$\dfrac{\pi^2 R^2\sigma_0 \mathrm{G}}{2\alpha}$	$= \dfrac{3\pi^2 R^2\sigma_0\mathrm{G}(4\alpha^2 - 3R^2)}{16\alpha^3}$
	$=$	$\dfrac{3\pi R^2 M\mathrm{G}}{4\alpha^3}$	$= \dfrac{15\pi R^2 M\mathrm{G}(4\alpha^2 - 3R^2)}{32\alpha^3}$
Disk edge velocity	$V_c^2(\alpha,0) =$	$\dfrac{\pi^2\alpha\sigma_0\mathrm{G}}{2}$	$= \dfrac{3\pi^2\alpha\sigma_0\mathrm{G}}{16}$
	$=$	$\dfrac{3\pi M\mathrm{G}}{4\alpha}$	$= \dfrac{15\pi M\mathrm{G}}{32\alpha}$

5. Example: The force field of a simple galaxy model

Three-dimensional problems such as that of the structure and kinematics of the extra-planar gas will benefit from the use of the new density-potential pairs. A simple galaxy model was constructed for illustration. The model consists of a an n=2 disk and a core/bulge region modeled as a point mass. This model is defined by three parameters: the mass of the disk, the mass of



Fig. 2.— Circular velocity, $V_c = \sqrt{(-VF_R(R,z)}$, of a simple galaxy model at z=0, z=2.6 kpc, and z=3.9 kpc above the disk. The model consists of an n=2 disk and a point mass to represent the core + bulge. The masses of the two components are $M_{disk} = 3.5 \times 10^{10}\mathrm{M}_{\odot}$ and $M_{PointMass} = 4 \times 10^{10}\mathrm{M}_{\odot}$. The radius of the disk is $\alpha = 19$ kpc.

the core/bulge region, and the diameter of the disk. As shown in figure 2, the circular velocity, calculated as $V_c = \sqrt{(-VF_R(R,z)}$, is nearly constant over much of the disk. Also, the derivative of the circular velocity with z is nearly linear over a wide range of both R and z.

Figure 2 agrees surprisingly well with figure 5 of Fraternali et al. (2005) which shows that the measured velocity of HI for NGC891 decreases linearly with the height above the disk. See also Rand (1997); Swaters et al. (1997); Kamphuis et al. (2007); Oosterloo et al. (2007); Fraternali & Binney (2006); Barnabe et al. (2006). Further work is planned on this topic.

6. Summary and conclusion

We have presented new solutions for a family of finite disks. Closed form expressions in cylindrical coordinates using elementary functions are given for the potential and gravitational force for the disks with surface density $\Sigma_n = \sigma_0(1 - R^2/\alpha^2)^{n-1/2}$ with $n = 0, 1, 2$. Expressions are also given for the limiting cases of $R = 0$ and $z = 0$.

These solutions fill a need and should make it easier to model 3-D gravitational phenomenon involving disk galaxies. This is particularly important due to the recent availability of detailed kinematic data above the plane of the disk.

I am grateful to the anonymous referee for useful suggestions and comments which improved the presentation.

A. Some simple identities

A number of identities are gathered here. In all cases $x, y \in \Re$ The ranges of validity must avoid the discontinuity of the principal value of the square root function on the negative real axis.

Equation A1 can be proved by taking the sin of both sides; reducing the terms using the identities $\sin(a+b) = \sin(a)\cos(b) + \cos(a)\sin(b)$ and $\sin(2a) = 2\sin(a)\cos(b)$; and substituting $\cos = \sqrt{1-\sin^2}$. The other identities can be found by substituting into the relation

$$\sqrt{x+\mathtt{I}\,y} = \frac{\sqrt{\sqrt{x^2+y^2}+x} + \mathrm{sgn}(y)\mathtt{I}\,\sqrt{\sqrt{x^2+y^2}-x}}{\sqrt{2}}$$

and into the expression found by taking the reciprocal of both sides.

$$\sin^{-1}(x-\mathtt{I}\,y) + \sin^{-1}(x+\mathtt{I}\,y) = 2\sin^{-1}\left[\tfrac{1}{2}\sqrt{(x+1)^2+y^2} - \tfrac{1}{2}\sqrt{(x-1)^2+y^2}\right] \tag{A1}$$

$$\sqrt{x-\mathtt{I}\,y} + \sqrt{x+\mathtt{I}\,y} = \sqrt{2}\sqrt{\sqrt{x^2+y^2}+x} \tag{A2}$$

$$\mathtt{I}\,\sqrt{x-\mathtt{I}\,y} - \mathtt{I}\,\sqrt{x+\mathtt{I}\,y} = \mathrm{sgn}(y)\sqrt{2}\sqrt{\sqrt{x^2+y^2}-x} \tag{A3}$$

$$\frac{1}{\sqrt{x+\mathtt{I}\,y}} + \frac{1}{\sqrt{x-\mathtt{I}\,y}} = \sqrt{2}\frac{\sqrt{\sqrt{x^2+y^2}+x}}{\sqrt{x^2+y^2}} \tag{A4}$$

$$\frac{\mathtt{I}}{\sqrt{x+\mathtt{I}\,y}} - \frac{\mathtt{I}}{\sqrt{x-\mathtt{I}\,y}} = \mathrm{sgn}(y)\sqrt{2}\frac{\sqrt{\sqrt{x^2+y^2}-x}}{\sqrt{x^2+y^2}} \tag{A5}$$

$$\sqrt{1-(x+\mathtt{I}\,y)^2} + \sqrt{1-(x-\mathtt{I}\,y)^2} = \sqrt{2}\sqrt{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}+1-x^2+y^2} \tag{A6}$$

$$\mathtt{I}\,\sqrt{1-(x+\mathtt{I}\,y)^2} - \mathtt{I}\,\sqrt{1-(x-\mathtt{I}\,y)^2} = \mathrm{sgn}(xy)\sqrt{2}\sqrt{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}-1+x^2-y^2} \tag{A7}$$

$$\frac{1}{\sqrt{1-(x-\mathtt{I}\,y)^2}} + \frac{1}{\sqrt{1-(x+\mathtt{I}\,y)^2}} = \sqrt{2}\frac{\sqrt{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}+1-x^2+y^2}}{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}} \tag{A8}$$

$$\frac{\mathtt{I}}{\sqrt{1-(x-\mathtt{I}\,y)^2}} - \frac{\mathtt{I}}{\sqrt{1-(x+\mathtt{I}\,y)^2}} = \mathrm{sgn}(xy)\sqrt{2}\frac{\sqrt{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}-1+x^2-y^2}}{\sqrt{1-2x^2+2y^2+x^4+2x^2y^2+y^4}} \tag{A9}$$

REFERENCES

Barnabe, M., Ciotti, L., Fraternali, F., & Sancisi, R. 2006, A&A, 446, 61

Bertin, G. 2000, Dynamics of galaxies (Cambridge University Press)

Binney, J. & Tremaine, S. 2008, Galactic Dynamics: Second Edition (Princeton University Press, Princeton, NJ.)

Casertano, S. 1983, MNRAS, 203, 735

Chandrasekhar, S. 1987, Ellipsoidal figures of equilibrium (New York : Dover, 1987)

Ciotti, L. & Giampieri, G. 2007, MNRAS, 376, 1162

Ciotti, L. & Marinacci, F. 2008, MNRAS, 387, 1117

Conway, J. T. 2000, MNRAS, 316, 540

Cuddeford, P. 1993, MNRAS, 262, 1076

Evans, N. W. & Collett, J. L. 1993, MNRAS, 264, 353

Evans, N. W. & de Zeeuw, P. T. 1992, MNRAS, 257, 152

Fraternali, F. & Binney, J. J. 2006, MNRAS, 366, 449

Fraternali, F., Oosterloo, T. A., Sancisi, R., & Swaters, R. 2005, in ASP Conf. Ser. 331: Extra-Planar Gas, 239–+

Freeman, K. C. 1970, ApJ, 160, 811

González, G. A. & Reina, J. I. 2006, MNRAS, 371, 1873

Gradshteyn, I. S. & Ryzhik, I. M. 1994, Table of integrals, series and products (Academic Press, 5th ed. Edited by Jeffrey, A)

Hunter, C. 1963, MNRAS, 126, 299

Hunter, J. H., Ball, R., & Gottesman, S. T. 1984, MNRAS, 208, 1

Huré, J.-M., Hersant, F., Carreau, C., & Busset, J.-P. 2008, A&A, 490, 477

Huré, J.-M. 2005, A&A, 434, 1

Kalnajs, A. J. 1971, ApJ, 166, 275

—. 1972, ApJ, 175, 63

Kamphuis, P., Peletier, R. F., Dettmar, R.-J., van der Hulst, J. M., van der Kruit, P. C., & Allen, R. J. 2007, A&A, 468, 951

Kuzmin, G. G. 1956, Astr.Zh., 33, 27

Lass, H. & Blitzer, L. 1983, Celestial Mechanics, 30, 225

Lynden-Bell, D. & Pineault, S. 1978, MNRAS, 185, 679

Lynden-Bell, D. 1989, MNRAS, 237, 1099

Meinel, R. 2001, in Exact Solutions and Scalar Fields in Gravity: Recent Developments (Edited by A Macias, J Cervantes-Cota, and C Lammerzahl; Kluwer Academic Publishers, New York), 69–75

Mestel, L. 1963, MNRAS, 126, 553

Mihalas, D. & Routly, P. M. 1968, Galactic astronomy (W.H. Freeman and Company)

Miyamoto, M. & Nagai, R. 1975, PASJ, 27, 533

Neugebauer, G. & Meinel, R. 1995, Physical Review Letters, 75, 3046

Oosterloo, T., Fraternali, F., & Sancisi, R. 2007, AJ, 134, 1019

Pedraza, J. F., Ramos-Caro, J., & Gonzalez, G. A. 2008, MNRAS, 390, 1587

Rand, R. J. 1997, ApJ, 474, 129

Satoh, C. 1980, PASJ, 32, 41

Schmidt, M. 1956, Bull. Astron. Inst. Netherlands, 13, 15

Swaters, R. A., Sancisi, R., & van der Hulst, J. M. 1997, ApJ, 491, 140

Toomre, A. 1963, ApJ, 138, 385

Vokrouhlicky, D. & Karas, V. 1998, MNRAS, 298, 53

This preprint was prepared with the AAS LaTeX macros v5.2.